



CHERT

Flint is a form of microcrystalline quartz that is typically called "chert" by geologists. Flint often forms as nodules in sedimentary rocks such as chalk and marine limestones. Some rock units form through the accumulation of siliceous skeletal material. These can recrystallize to form a layer of bedded flint. Flint is highly resistant to weathering. Early people who used flint to make tools often found nicely shaped pieces of this for making specific tools.



CHERT

Flint is a form of microcrystalline quartz that is typically called "chert" by geologists. Flint often forms as nodules in sedimentary rocks such as chalk and marine limestones. Some rock units form through the accumulation of siliceous skeletal material. These can recrystallize to form a layer of bedded flint. Flint is highly resistant to weathering. Early people who used flint to make tools often found nicely shaped pieces of this for making specific tools.



CHERT

Flint is a form of microcrystalline quartz that is typically called "chert" by geologists. Flint often forms as nodules in sedimentary rocks such as chalk and marine limestones. Some rock units form through the accumulation of siliceous skeletal material. These can recrystallize to form a layer of bedded flint. Flint is highly resistant to weathering. Early people who used flint to make tools often found nicely shaped pieces of this for making specific tools.



CHERT

Flint is a form of microcrystalline quartz that is typically called "chert" by geologists. Flint often forms as nodules in sedimentary rocks such as chalk and marine limestones. Some rock units form through the accumulation of siliceous skeletal material. These can recrystallize to form a layer of bedded flint. Flint is highly resistant to weathering. Early people who used flint to make tools often found nicely shaped pieces of this for making specific tools.



CHERT

Flint is a form of microcrystalline quartz that is typically called "chert" by geologists. Flint often forms as nodules in sedimentary rocks such as chalk and marine limestones. Some rock units form through the accumulation of siliceous skeletal material. These can recrystallize to form a layer of bedded flint. Flint is highly resistant to weathering. Early people who used flint to make tools often found nicely shaped pieces of this for making specific tools.



CHERT

Flint is a form of microcrystalline quartz that is typically called "chert" by geologists. Flint often forms as nodules in sedimentary rocks such as chalk and marine limestones. Some rock units form through the accumulation of siliceous skeletal material. These can recrystallize to form a layer of bedded flint. Flint is highly resistant to weathering. Early people who used flint to make tools often found nicely shaped pieces of this for making specific tools.