



CRINOID

Stalked Crinoids, or “sea lilies”, lived in shallow marine habitats during the Paleozoic era until the Permo-Triassic extinction. They attached themselves to the ocean floor, and filtered food particles from the currents flowing past them. Many Paleozoic limestones are made up largely of crinoid skeletal fragments.



CRINOID

Stalked Crinoids, or “sea lilies”, lived in shallow marine habitats during the Paleozoic era until the Permo-Triassic extinction. They attached themselves to the ocean floor, and filtered food particles from the currents flowing past them. Many Paleozoic limestones are made up largely of crinoid skeletal fragments.



CRINOID

Stalked Crinoids, or “sea lilies”, lived in shallow marine habitats during the Paleozoic era until the Permo-Triassic extinction. They attached themselves to the ocean floor, and filtered food particles from the currents flowing past them. Many Paleozoic limestones are made up largely of crinoid skeletal fragments.



CRINOID

Stalked Crinoids, or “sea lilies”, lived in shallow marine habitats during the Paleozoic era until the Permo-Triassic extinction. They attached themselves to the ocean floor, and filtered food particles from the currents flowing past them. Many Paleozoic limestones are made up largely of crinoid skeletal fragments.



CRINOID

Stalked Crinoids, or “sea lilies”, lived in shallow marine habitats during the Paleozoic era until the Permo-Triassic extinction. They attached themselves to the ocean floor, and filtered food particles from the currents flowing past them. Many Paleozoic limestones are made up largely of crinoid skeletal fragments.



CRINOID

Stalked Crinoids, or “sea lilies”, lived in shallow marine habitats during the Paleozoic era until the Permo-Triassic extinction. They attached themselves to the ocean floor, and filtered food particles from the currents flowing past them. Many Paleozoic limestones are made up largely of crinoid skeletal fragments.