



Brittle Starfish Fossil

Brittle starfish or ophiuroids are echinoderms and are closely related to the starfish. These specimens were found in the Sahara Desert region of Morocco. The fossil is from the Ordovician Period around 488 to 443 million years ago.



Brittle Starfish Fossil

Brittle starfish or ophiuroids are echinoderms and are closely related to the starfish. These specimens were found in the Sahara Desert region of Morocco. The fossil is from the Ordovician Period around 488 to 443 million years ago.



Brittle Starfish Fossil

Brittle starfish or ophiuroids are echinoderms and are closely related to the starfish. These specimens were found in the Sahara Desert region of Morocco. The fossil is from the Ordovician Period around 488 to 443 million years ago.



Brittle Starfish Fossil

Brittle starfish or ophiuroids are echinoderms and are closely related to the starfish. These specimens were found in the Sahara Desert region of Morocco. The fossil is from the Ordovician Period around 488 to 443 million years ago.



Brittle Starfish Fossil

Brittle starfish or ophiuroids are echinoderms and are closely related to the starfish. These specimens were found in the Sahara Desert region of Morocco. The fossil is from the Ordovician Period around 488 to 443 million years ago.



Brittle Starfish Fossil

Brittle starfish or ophiuroids are echinoderms and are closely related to the starfish. These specimens were found in the Sahara Desert region of Morocco. The fossil is from the Ordovician Period around 488 to 443 million years ago.